

$\nu_\mu \leftrightarrow \nu_\tau$ vs $\nu_\mu \leftrightarrow \nu_s$ solutions for the atmospheric neutrino problem *

Osamu Yasuda

Department of Physics, Tokyo Metropolitan University
1-1 Minami-Osawa Hachioji, Tokyo 192-0397, Japan

The $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ solutions to the atmospheric neutrino problem are compared with Superkamiokande data. Both the solutions with a large mixing angle seem to be consistent with the data.

1. Introduction

Recent atmospheric neutrino data by Superkamiokande [1–3] provide strong evidence for neutrino oscillations. It has been shown [1–3] that atmospheric neutrino data favor $\nu_\mu \leftrightarrow \nu_\tau$ oscillations with maximal mixing, rather than $\nu_\mu \leftrightarrow \nu_e$. However, ν_μ disappearance alone does not imply uniquely a $\nu_\mu \leftrightarrow \nu_\tau$ solution and there is another solution $\nu_\mu \leftrightarrow \nu_s$, where ν_s denotes a sterile neutrino. In this talk some aspects of the $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ solutions are discussed.

In the past there has been a prejudice against the $\nu_\mu \leftrightarrow \nu_s$ solution to the atmospheric neutrino problem. The argument [4] was based on big bang nucleosynthesis which gives a condition $\Delta m^2 \sin^4 2\theta \lesssim 10^{-4} \text{eV}^2$ in order for sterile neutrinos not to be in thermal equilibrium. However, there was a loophole in this argument. Foot and Volkas [5] have shown that large lepton asymmetries will suppress $\nu_s \leftrightarrow \nu_x$ neutrino oscillations. Interestingly, given certain conditions, the required lepton asymmetries can actually be created by the oscillations themselves [5]. So there is no longer any obstruction to $\nu_\mu \leftrightarrow \nu_s$ as a solution to the atmospheric neutrino anomaly.

Many models [6,7] have been proposed which predict large or maximal active-sterile mixing. Among others, Foot and Volkas have been observed by exact parity symmetric models [7] and this was the main motivation of [8] in which $\nu_\mu \leftrightarrow \nu_s$ was examined in detail by fitting to the contained events of the Superkamiokande atmo-

spheric neutrino data for 414 days.

2. Analysis of the Superkamiokande contained events for 414 days

The survival probability $P(\nu_\alpha \leftrightarrow \nu_\alpha)$ is obtained by solving the Schrödinger equation for neutrino evolution including matter effects. It is given by

$$i \frac{d}{dx} \begin{pmatrix} \nu_\mu(x) \\ \nu_{\tau,s}(x) \end{pmatrix} = M \begin{pmatrix} \nu_\mu(x) \\ \nu_{\tau,s}(x) \end{pmatrix}, \quad (1)$$

$$M \equiv U \text{diag}(0, \frac{\Delta m^2}{2E}) U^{-1} + \text{diag}(0, A_{\tau,s}(x)),$$

where

$$U \equiv \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

is the MNS mixing matrix [9], x is the distance traveled, Δm^2 the difference in squared masses, θ the vacuum mixing angle and $\nu_{\mu,\tau,s}(x)$ the wavefunctions of the neutrinos. The quantities $A_{\tau,s}(x)$ are the effective potential differences generated through the matter effect [10]:

$$A_\tau(x) = 0$$

and, for electrically neutral terrestrial matter [11]

$$A_s(x) = \frac{1}{\sqrt{2}} G_F N_n(x),$$

where G_F is the Fermi constant, $N_n(x)$ is the number density of neutrons along the path of the neutrino. It is this matter effect $A_{\tau,s}$ that make a difference between the $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ oscillations. For antineutrinos the sign of A_s is reversed.

*Talk presented at “NEUTRINO 98”, Takayama, Japan, June 4-9, 1998.

The way to obtain the numbers of events and evaluate χ^2 is described in [8], where two quantities have been introduced to perform a χ^2 analysis. One is the double ratio [12]

$$R \equiv \frac{(N_\mu/N_e)|_{\text{osc}}}{(N_\mu/N_e)|_{\text{no-osc}}}$$

where the quantities $N_{e,\mu}$ are the numbers of e -like and μ -like events. The numerator denotes numbers with oscillation probability obtained by (2), while the denominator the numbers expected with oscillations switched off. The other one is the quantity on up-down flux asymmetries for α -like ($\alpha=e,\mu$) events and is defined by

$$Y_\alpha \equiv \frac{(N_\alpha(\cos \Theta < -0.2)/N_\alpha(\cos \Theta > 0.2))|_{\text{osc}}}{(N_\alpha(\cos \Theta < -0.2)/N_\alpha(\cos \Theta > 0.2))|_{\text{no-osc}}},$$

where Θ is the zenith angle, $N_\alpha(\cos \Theta < -0.2)$ and $N_\alpha(\cos \Theta > 0.2)$ are the number of upward and downward going events, respectively. χ^2 with the double ratio R is defined by

$$\chi_{\text{atm}}^2(R) = \sum_E \left(\frac{R^{SK} - R^{th}}{\delta R^{SK}} \right)^2,$$

and χ^2 with the up-down asymmetry Y_α is defined by

$$\begin{aligned} & \chi_{\text{atm}}^2(Y) \\ &= \sum_E \left[\left(\frac{Y_\mu^{SK} - Y_\mu^{th}}{\delta Y_\mu^{SK}} \right)^2 + \left(\frac{Y_e^{SK} - Y_e^{th}}{\delta Y_e^{SK}} \right)^2 \right], \end{aligned}$$

where the sum is over the sub-GeV and multi-GeV cases, the measured Superkamiokande values and errors are denoted by the superscript “SK” and the theoretical predictions for the quantities are labeled by “th”.

The results of the χ^2 fits are displayed in Figs.1–8. In Figs. 1 and 2, χ^2 is plotted against Δm^2 . For $\nu_\mu \leftrightarrow \nu_\tau$, χ^2 does not experience a deep minimum at the best fit point with respect to Δm^2 particularly when the R ’s are excluded from the fit. In general, for geometrical reasons, atmospheric neutrino analysis does not constrain Δm^2 very precisely. Note that the the situation is slightly different in case of $\nu_\mu \leftrightarrow \nu_s$. Figure 3 shows the allowed region of $(\sin^2 2\theta, \Delta m^2)$ at




Figure 1. χ^2 as a function of Δm^2 .

- (a): $\nu_\mu \leftrightarrow \nu_\tau$, $\sin^2 2\theta = 1$;
- (b): $\nu_\mu \leftrightarrow \nu_\tau$, $\sin^2 2\theta = 0.8$;
- (c): $\nu_\mu \leftrightarrow \nu_s$, $\sin^2 2\theta = 1$;
- (d): $\nu_\mu \leftrightarrow \nu_s$, $\sin^2 2\theta = 0.8$, $\Delta m^2 > 0$;
- (e): $\nu_\mu \leftrightarrow \nu_s$, $\sin^2 2\theta = 0.8$, $\Delta m^2 < 0$.




Figure 2. The same as Figure 1 but with R data excluded from the fit.




Figure 3. The allowed region in the $(\sin^2 2\theta, \Delta m^2)$ plane for the $\nu_\mu \leftrightarrow \nu_\tau$ scenario.




Figure 5. The allowed region in the $(\sin^2 2\theta, \Delta m^2)$ plane for the $\nu_\mu \leftrightarrow \nu_s$ scenario with $\Delta m^2 > 0$.




Figure 4. The same as Figure 3 but with R data excluded from the fit.




Figure 6. The same as Figure 5 but with R data excluded from the fit.




Figure 7. The allowed region in the $(\sin^2 2\theta, \Delta m^2)$ plane for the $\nu_\mu \leftrightarrow \nu_s$ scenario with $\Delta m^2 < 0$.




Figure 8. The same as Figure 7 but with R data excluded from the fit.

various confidence levels for the $\nu_\mu \leftrightarrow \nu_\tau$ scenario. Maximal mixing provides the best fit, and Δm^2 values in the 10^{-3} to 10^{-2} eV² range are favored. Note that the confidence levels are defined in the usual way by

$$\chi^2 = \chi_{\min}^2 + \Delta\chi^2$$

where $\Delta\chi^2 = 2.3, 4.6, 6.2, 11.8$ for the 1 σ , 90% C.L., 2 σ and 3 σ allowed region respectively. Our $\chi_{\min}^2$ for $\nu_\mu \leftrightarrow \nu_\tau$ oscillations is $\chi_{\min}^2 = 4.5$ for 4 degrees of freedom. This is quite a good fit to the data (allowed at the 35% level). In Figure 4 we show the allowed region considering just the asymmetries instead of using both the asymmetries and the R ratios. Note that in this case there are 4 data points and 2 free parameters which gives 2 degrees of freedom.

Figures 5–8 show the corresponding results for the $\nu_\mu \leftrightarrow \nu_s$ scenario. If $\Delta m^2 > 0$, smaller values of Δm^2 are disfavored because the matter effect moves both R and Y away from the measured values, but if $\Delta m^2 < 0$, then smaller values of Δm^2 and $\sin^2 2\theta$ are permitted at the 90% confidence level. The value of $\chi_{\min}^2$ for the $\nu_\mu \leftrightarrow \nu_s$ scenario is $\chi_{\min}^2 = 5.1$ for 4 degrees of freedom. This is similar to $\nu_\mu - \nu_\tau$ case and also represents quite a good fit (which is allowed at 28%).

To summarize, both the solutions $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ provide a good fit to the contained events of the Superkamiokande atmospheric neutrino data.

3. Other analyses

There have been several proposals to distinguish the $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ oscillations. Matter effects in $\nu_\mu \leftrightarrow \nu_s$ oscillations in upward going muon data were first analyzed by Akhmedov, Lipari and Lusignoli [13] and more recently by Lipari and Lusignoli [14]. It has been pointed out by Liu, Smirnov [15] and Liu, Mikheyev, Smirnov [16] that signatures due to parametric enhancement in $\nu_\mu \leftrightarrow \nu_s$ oscillations may be seen in upward going muon data. Vissani and Smirnov [17] proposed to look at the ratio $(\pi^0\text{-events})/(\text{two ring events})$. Learned, Pakvasa and Stone [18] suggested that the up-down asymmetry $(\text{upward going } \pi^0\text{-events})/(\text{downward going } \pi^0\text{-events})$ can

tell a difference. Hall and Murayama [19] proposed a similar technique to use the up-down asymmetry in the multi-ring events. Kajita [2] mentioned the ratio (π^0 -events)/(e-like events) which should in principle enable us to distinguish. All these analyses seem to be still inconclusive and we need more statistics and accurate knowledge on nuclear cross sections to draw a conclusion.

4. Conclusions

We have demonstrated that matter effects in the Earth have a significant role to play in comparing and contrasting the $\nu_\mu \leftrightarrow \nu_\tau$ and $\nu_\mu \leftrightarrow \nu_s$ solutions to the atmospheric neutrino anomaly with Superkamiokande data. So far both solutions provide a good fit to the data and we need more statistics to be conclusive. We hope that non-accelerator experiments such as Superkamiokande will distinguish them before future long baseline experiments with emulsion techniques [20] give direct evidence.

Acknowledgments

The author would like to thank R. Foot and R.R. Volkas for collaboration, discussions and comments on this manuscript. He also would like to thank T. Kajita and E. Kearns for useful communications. This research was supported in part by a Grant-in-Aid for Scientific Research of the Ministry of Education, Science and Culture, #09045036, #10140221, #10640280.

REFERENCES

1. Y. Fukuda et al., Phys. Lett. **B433** (1998) 9; hep-ex/9805006.
2. T. Kajita, these proceeding; <http://www-sk.icrr.u-tokyo.ac.jp/nu98/scan/063/>.
3. Y. Fukuda et al., hep-ex/9807003.
4. R. Barbieri and A. Dolgov, Phys. Lett. **B237** (1990) 440, Nucl. Phys. **B349** (1991) 743; K. Kainulainen, Phys. Lett. **B244** (1990) 191; K. Enqvist, K. Kainulainen and M. Thomson, Nucl. Phys. **B373** (1992) 498, Phys. Lett. **B288** (1992) 145; X. Shi, D.N. Schramm and B.D. Fields, Phys. Rev. **D48** (1993) 2563.
5. R. Foot and R. R. Volkas, Phys. Rev. **D55** (1997) 5147; Astropart. Phys. **7** (1997) 283; Phys. Rev. **D56** (1997) 6653.
6. M. Kobayashi, C. S. Lim and M. M. Nojiri, Phys. Rev. Lett. **67** (1991) 1685; C. Giunti, C. W. Kim and U. W. Lee, Phys. Rev. **D46** (1992) 3034; J. Bowes and R. R. Volkas, J. Phys. **G24** (1998) 1249; A. Geiser, preprint CERN-EP-98-056.
7. R. Foot, Mod. Phys. Lett. **A9** (1994) 169; R. Foot and R. R. Volkas, Phys. Rev. **D52** (1995) 6595.
8. R. Foot, R. R. Volkas and O. Yasuda, Phys. Rev. **D58** (1998) 13006.
9. Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
10. L. Wolfenstein, Phys. Rev. **D17** (1978) 2369; S.P. Mikheyev and A.Yu. Smirnov, Yad. Fiz. **42** (1985) 1441 [Sov. J. Nucl. Phys. **42** (1985) 913]; Nuovo Cim. **C9** (1986) 17.
11. See for example, D. Notzold and G. Raffelt, Nucl. Phys. **B307** (1988) 924.
12. K.S. Hirata et al., Phys. Lett. **B280** (1992) 146; Y. Fukuda et al., Phys. Lett. **B335** (1994) 237.
13. E. Akhmedov, P. Lipari and M. Lusignoli, Phys. Lett. **B300** (1993) 128.
14. P. Lipari and M. Lusignoli, hep-ph/9803440.
15. Q.Y. Liu and A.Yu. Smirnov, hep-ph/9712493.
16. Q.Y. Liu, S.P. Mikheyev and A.Yu. Smirnov, hep-ph/9803415.
17. F. Vissani and A.Yu. Smirnov, Phys. Lett. **B432** (1998) 376; See also T. Kajita, Talk at *Topical Workshop on Neutrino Physics*, Institute for Theoretical Physics, The University of Adelaide, Nov. 1996.
18. J.G. Learned, S. Pakvasa and J.L. Stone, hep-ph/9805343. The author has learned from E. Kearns that this technique has been thought about for some time by the members of the Superkamiokande collaboration.
19. L. Hall and H. Murayama, hep-ph/9806218.
20. MINOS experiment, <http://www.hep.anl.gov/NDK/HyperText/numi.html>; OPERA experiment, <http://www1.na.infn.it/wsbsubnucl/accel/neutrino/opera.html>; ICARUS experiment, <http://www.aquila.infn.it/icarus/>.